

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	11.5 (8.8-15.0)	799	20.4 (12.1-32.2)	88			●
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	19.0 (16.7-21.6)	800	32.6 (21.9-45.5)	88		●	
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	5.2 (3.3-8.3)	500	32.5 (22.9-43.7)	45		●	
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	43.0 (38.8-47.3)	512	46.9 (35.7-58.4)	49			●
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	23.5 (19.6-28.0)	774	29.1 (20.1-40.2)	82			●
QN14: Percentage of students who carried a gun on one or more of the past 30 days	8.1 (5.9-11.1)	789	16.4 (11.1-23.6)	86		●	
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	7.1 (5.2-9.7)	796	11.6 (6.6-19.5)	82			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	7.5 (5.9-9.5)	783	18.0 (11.8-26.6)	85		●	
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	5.3 (3.8-7.3)	806	18.5 (11.4-28.6)	88		●	
QN18: Percentage of students who were in a physical fight one or more times during the past 12 months	26.6 (22.9-30.7)	791	41.3 (31.7-51.6)	83		●	
QN19: Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse	3.0 (2.0-4.4)	791	13.8 (7.6-23.8)	85		●	
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	7.3 (4.9-10.9)	795	13.8 (7.8-23.3)	85			●
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	11.5 (9.4-14.1)	800	33.0 (24.2-43.3)	87		●	
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	10.6 (8.0-13.9)	677	29.6 (18.9-43.2)	75		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN23: Among students who dated or went out with someone during the past 12 months, the percentage who had been forced by someone they were dating or going out with to do sexual things that they did not want to one or more times during the past 12 months	11.6 (9.4-14.2)	682	21.3 (15.4-28.6)	74		●	
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	20.0 (16.8-23.6)	804	42.7 (33.4-52.6)	87		●	
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	13.6 (11.2-16.5)	802	42.4 (30.8-54.8)	87		●	
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	29.3 (25.9-32.9)	805	62.3 (50.7-72.6)	86		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	17.4 (13.6-22.1)	804	43.6 (34.3-53.3)	87		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	13.1 (10.5-16.2)	803	45.6 (33.5-58.2)	86		●	
QN29: Percentage of students who actually attempted suicide one or more times during the past 12 months	8.2 (6.1-10.8)	691	33.0 (24.3-43.1)	75		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [‡]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [‡]	No Difference [‡]
	% 95% CI [*]	N	% 95% CI	N			
QN30: Percentage of students who made a suicide attempt during the past 12 months that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse	3.5 (2.3-5.2)	690	10.9 (5.3-21.0)	75		●	

*Confidence interval.

[‡]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who ever tried cigarette smoking, even one or two puffs	61.6 (55.6-67.2)	770	72.2 (58.5-82.7)	78			●
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	17.2 (13.9-21.1)	767	35.0 (24.1-47.7)	78		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	24.8 (20.8-29.2)	746	37.8 (25.5-51.9)	79			●
QNFRCIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	10.0 (7.8-12.8)	746	18.7 (10.4-31.4)	79			●
QN34: Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days	10.9 (7.1-16.3)	181	19.5 (10.4-33.8)	30			●
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	25.2 (18.5-33.5)	157	-	21			
QN36: Percentage of students who smoked cigarettes on school property on one or more of the past 30 days	6.5 (4.6-9.1)	785	18.4 (10.8-29.7)	83		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN37: Percentage of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days	18.6 (15.5-22.1)	772	27.4 (18.1-39.1)	79			●
QN38: Among students who reported current cigarette use, the percentage who did not ever try to quit smoking cigarettes during the past 12 months	40.3 (30.4-51.0)	179	46.9 (27.2-67.6)	30			●
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	18.6 (14.4-23.7)	806	16.8 (9.0-29.2)	88			●
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	17.1 (14.6-20.0)	804	30.3 (18.5-45.3)	87		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	36.9 (31.7-42.4)	745	43.7 (28.3-60.5)	78			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	79.1 (75.2-82.5)	775	85.2 (75.2-91.6)	82			●
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	21.5 (17.2-26.4)	802	31.9 (20.8-45.5)	87			●
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	43.3 (39.2-47.4)	706	58.3 (43.1-72.2)	79		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	28.1 (24.3-32.2)	755	42.5 (28.4-58.0)	81			●
QN45: Percentage of students who had ten or more drinks of alcohol in a row, that is, within a couple of hours, during the past 30 days	9.2 (7.1-12.0)	721	16.4 (9.4-27.1)	80			●
QN46: Among students who reported current alcohol use, the percentage who usually got the alcohol they drank from someone who gave it to them during the past 30 days	35.1 (30.3-40.2)	303	26.7 (15.9-41.2)	46			●
QN47: Percentage of students who used marijuana one or more times during their life	49.4 (44.2-54.6)	774	56.8 (45.0-67.9)	84			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN48: Percentage of students who tried marijuana for the first time before age 13 years	11.6 (8.8-15.2)	789	17.9 (10.5-28.7)	84			●
QN49: Percentage of students who used marijuana one or more times during the past 30 days	25.3 (21.1-30.1)	794	40.6 (29.3-53.0)	86		●	
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	5.7 (4.1-8.0)	803	14.8 (8.2-25.2)	86		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	7.8 (6.1-10.0)	804	26.4 (17.2-38.4)	88		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	3.2 (2.0-5.0)	804	16.0 (9.2-26.3)	88		●	
QN54: Percentage of students who used ecstasy one or more times during their life	4.8 (3.4-6.6)	803	18.0 (9.9-30.3)	87		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	2.5 (1.4-4.3)	805	15.5 (10.6-22.2)	88		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	15.5 (12.5-19.0)	804	41.8 (30.7-53.9)	88		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	25.2 (22.8-27.8)	802	27.8 (17.7-40.8)	85			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	76.5 (71.5-80.8)	724	79.0 (68.0-87.0)	76			●
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	8.5 (6.2-11.6)	723	13.6 (6.1-27.5)	76			●
QN61: Percentage of students who had sexual intercourse with four or more people during their life	20.2 (16.6-24.3)	718	31.9 (19.9-46.9)	75			●
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	54.1 (48.0-60.0)	718	60.8 (45.5-74.2)	75			●
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	16.1 (12.5-20.6)	381	33.9 (20.7-50.2)	43		●	
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	45.1 (40.1-50.2)	377	57.5 (40.2-73.1)	42			●
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	79.3 (74.4-83.4)	376	89.3 (75.4-95.8)	42			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	92.0 (88.0-94.8)	376	97.5 (83.2-99.7)	42			●
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	71.3 (66.0-76.0)	376	86.8 (72.8-94.1)	42		●	
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	93.1 (89.9-95.3)	372	95.6 (84.9-98.8)	41			●
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	13.5 (10.2-17.6)	376	28.4 (16.1-45.1)	42			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [‡]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [‡]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) [§]	16.9 (14.4-19.7)	777	10.8 (5.6-19.9)	82			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) [§]	16.4 (13.5-19.9)	777	21.3 (13.2-32.6)	82			●
QN66: Percentage of students who described themselves as slightly or very overweight	28.9 (25.4-32.7)	795	49.4 (37.3-61.6)	88		●	
QN68: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days	14.3 (11.3-17.9)	794	33.8 (20.9-49.8)	83		●	
QN69: Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days	7.8 (6.0-10.1)	795	22.5 (12.9-36.3)	82		●	
QN70: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days	5.9 (4.4-7.8)	796	14.7 (9.7-21.6)	82		●	
QN71: Percentage of students who did not drink 100% fruit juices during the past seven days	24.9 (21.7-28.4)	803	30.5 (18.0-46.6)	88			●

*Confidence interval.

[‡]Based on t-test analyses, p<0.05.

[§]Based on reference data from the 2000 CDC Growth Charts.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN72: Percentage of students who did not eat fruit during the past seven days	12.8 (10.6-15.5)	801	12.4 (6.8-21.7)	86			●
QN73: Percentage of students who did not eat green salad during the past seven days	46.7 (41.6-51.8)	803	48.7 (39.7-57.8)	87			●
QN74: Percentage of students who did not eat potatoes during the past seven days	24.3 (21.1-27.8)	800	20.4 (11.5-33.5)	86			●
QN75: Percentage of students who did not eat carrots during the past seven days	56.9 (51.7-61.9)	793	54.1 (43.6-64.3)	86			●
QN76: Percentage of students who did not eat other vegetables during the past seven days	20.3 (16.6-24.6)	799	18.5 (9.6-32.7)	88			●
QNFRVG: Percentage of students who ate fruits or drank 100% fruit juices or ate vegetables less than five times per day during the past seven days	83.2 (79.8-86.1)	788	82.7 (71.9-90.0)	84			●
QNFRUIT: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day during the past seven days	73.1 (68.1-77.5)	800	74.7 (61.1-84.8)	86			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who ate fruits or drank 100% fruit juices 0 times per day during the past seven days	6.6 (5.1-8.7)	800	8.6 (4.2-16.7)	86			●
QNFR1: Percentage of students who ate fruits or drank 100% fruit juices less than one time per day during the past seven days	41.4 (36.9-46.1)	800	48.3 (35.3-61.6)	86			●
QNFR3: Percentage of students who ate fruits or drank 100% fruit juices less than three times per day during the past seven days	83.1 (79.9-86.0)	800	81.5 (69.3-89.6)	86			●
QNVEG: Percentage of students who ate vegetables less than three times per day during the past seven days	87.4 (84.9-89.6)	791	84.8 (76.2-90.7)	85			●
QNVEG0: Percentage of students who ate vegetables 0 times per day during the past seven days	5.1 (3.5-7.5)	791	6.8 (2.3-18.2)	85			●
QNVEG1: Percentage of students who ate vegetables less than one time per day during the past seven days	41.4 (37.7-45.2)	791	35.8 (25.4-47.7)	85			●
QNVEG2: Percentage of students who ate vegetables less than two times per day during the past seven days	76.5 (72.8-79.8)	791	77.2 (68.0-84.4)	85			●
QNFRVG2: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day and ate vegetables less than three times per day during the past seven days	91.5 (89.0-93.5)	788	90.9 (83.0-95.3)	84			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	36.1 (30.8-41.6)	797	38.9 (28.4-50.5)	87			●
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	81.6 (77.1-85.3)	797	79.9 (71.0-86.5)	87			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	26.7 (22.3-31.7)	797	25.2 (15.2-38.7)	87			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	17.9 (14.2-22.3)	797	16.2 (9.5-26.2)	87			●
QN78: Percentage of students who did not drink milk or who drank less than three glasses per day of milk during the past seven days	90.6 (87.8-92.8)	793	87.5 (80.5-92.3)	87			●
QNMILK0: Percentage of students who drank 0 glasses per day of milk during the past seven days	21.3 (18.3-24.6)	793	25.9 (18.3-35.3)	87			●
QNMILK1: Percentage of students who drank less than one glass per day of milk during the past seven days	66.7 (62.6-70.5)	793	60.8 (53.6-67.6)	87			●
QNMILK2: Percentage of students who drank less than two glasses per day of milk during the past seven days	81.0 (77.2-84.2)	793	74.1 (66.8-80.3)	87			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI [*]	N	% 95% CI	N			
QN79: Percentage of students who did not eat breakfast on all of the past seven days	62.5 (58.8-66.1)	792	68.6 (58.6-77.2)	87			●
QNN0BKFT: Percentage of students who ate breakfast on none of the past seven days	12.1 (9.5-15.3)	792	22.8 (16.0-31.4)	87		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	57.2 (52.9-61.3)	789	71.3 (60.3-80.2)	88		●	
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	17.6 (14.4-21.4)	789	29.0 (22.0-37.2)	88		●	
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	74.4 (70.3-78.2)	789	82.8 (73.2-89.5)	88			●
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	27.7 (23.8-32.0)	795	20.4 (14.3-28.3)	88	●		
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	32.9 (29.6-36.4)	798	39.2 (28.3-51.2)	87			●
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	64.8 (56.1-72.6)	789	62.9 (48.8-75.1)	86			●
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	79.9 (74.3-84.5)	789	87.2 (75.1-93.9)	86			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [‡]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [‡]	No Difference [‡]
	% 95% CI [*]	N	% 95% CI	N			
QN84: Percentage of students who did not play on sports teams during the past 12 months	48.3 (43.6-53.0)	796	66.1 (56.0-74.9)	85		●	

*Confidence interval.

[‡]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Risk Behaviors and Sex of Sexual Partners Report

Other	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
Health Risk Behavior							
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	19.8 (16.1-24.0)	795	21.7 (15.0-30.4)	84			●
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	23.5 (20.9-26.2)	783	28.6 (21.0-37.7)	85			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who would not describe their grades in school as mostly A's or mostly B's during the past 12 months	32.2 (26.7-38.3)	798	40.1 (29.5-51.7)	87			●
QN88: Percentage of students who had seen or heard commercials on TV, the Internet, or on the radio about the dangers of cigarette smoking on two or less of the past 30 days	48.7 (45.5-51.8)	798	53.4 (41.7-64.8)	87			●
QN89: Percentage of students who reported that smoking was allowed only at some times or in some places or always allowed inside the home where they live	32.3 (28.1-36.8)	794	47.4 (34.5-60.6)	86		●	
QN90: Percentage of students who got the prescription drugs they took without a doctor's prescription by buying them from a family member or a friend; by taking them from a family member or a friend; or were given to them by a family member or a friend	11.5 (9.1-14.5)	798	33.0 (23.8-43.7)	87		●	
QN91: Percentage of students who have used synthetic marijuana (also called K2 or Spice) one or more times during their life	20.5 (17.7-23.6)	805	37.1 (24.7-51.5)	88		●	
QN92: Percentage of students who have ever had oral sex	66.4 (62.4-70.2)	793	86.5 (77.2-92.4)	84		●	
QN94: Percentage of students who drank a can, bottle, or glass of an energy drink, such as Red Bull or Jolt, one or more times per day during the past seven days	8.0 (6.1-10.5)	793	11.0 (5.1-22.0)	87			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Risk Behaviors and Sex of Sexual Partners Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN95: Percentage of students who have not ever talked or are not sure if they have ever talked about AIDS or HIV infection with their parents or other adults in their family	51.7 (47.5-55.8)	786	46.7 (34.4-59.5)	82			●
QN96: Percentage of students who brushed their teeth on six or less of the past seven days	21.3 (18.2-24.9)	795	36.9 (26.2-49.0)	85		●	
QN97: Percentage of students who flossed their teeth on six or less of the past seven days	82.3 (79.2-84.9)	797	86.7 (76.9-92.8)	86			●
QN98: Percentage of students who never saw, who last saw more than 12 months ago, or who were not sure when they last saw, a dentist for a check-up, exam, teeth cleaning, or other dental work	32.8 (28.5-37.4)	796	45.8 (33.2-59.1)	86			●
QN99: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems	16.5 (13.9-19.5)	782	34.8 (24.6-46.6)	84		●	
QN100: Percentage of students who definitely will not, probably will not, or are not sure if they will complete a post high school program such as a vocational training program, military service, community college, or 4-year college	17.8 (14.5-21.6)	791	24.1 (16.7-33.6)	85			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.